

200 mg of Zen

*Natural Support for Awake Calmness**

(Hypoallergenic)



Item # 74700

60 Vegetarian Capsules

The Possible Benefits of 200 mg of Zen, a Dietary Supplement

- May help support healthy emotional status*
 - May support a feeling of general relaxation without sedation*
 - May promote healthy moods during times of hormonal change*
-

Description

200 mg of Zen contains a significant quantity of both gamma-aminobutyric acid (GABA) and theanine (glutamic acid gamma-ethylamide), an amino acid derivative found naturally in green tea (*Camellia sinensis*). These two dietary supplement ingredients may together support healthy moods and a feeling of alert relaxation.*

Research has shown that dietary components can modulate the body's levels of neurotransmitters, e.g. serotonin, dopamine, norepinephrine, and GABA. Alterations in the levels of these neurotransmitters, perhaps induced by metabolic stress or vitamin deficiencies, e.g. vitamin B6, can significantly influence mood and emotional status, as well as motor function.*

GABA is the major inhibitory neurotransmitter in the brain and is active at 20% of central nervous system (CNS) synapses. GABA, via its neuronal A receptor, inhibits neurons by causing an influx of chloride ions. This chloride influx initiated by GABA

is known to be part of the bodily mechanisms involved with mood, muscle relaxation and sedation.* A decrease in GABA's function as an inhibitory mediator or a dysregulated sensitivity of GABA receptors can lead to excessive neuronal activity.*

Theanine, an amino acid derivative found naturally in green tea (*Camellia sinensis*), is also recognized to have calming properties.* Human studies have demonstrated that dietary theanine supplementation increases alpha wave activity, fostering a state of alert relaxation.* Animal studies have also shown that administration of theanine regulates brain serotonin concentration by affecting either serotonin synthesis or degradation in the brain.* It has been hypothesized that serotonin facilitates general and conditioned anxiety at the level of medial temporal lobe structures.

The calming effect of this tea component may seem contradictory to the stimulatory properties of green tea's caffeine content, however research

*THESE STATEMENTS HAVE NOT BEEN EVALUATED BY THE FOOD AND DRUG ADMINISTRATION. THIS PRODUCT IS NOT INTENDED TO DIAGNOSE, TREAT, CURE, OR PREVENT ANY DISEASE.

suggests that theanine exerts an antagonistic effect on caffeine's stimulatory action.* Although L-theanine is a component of green tea,

200 mg of Zen does not contain caffeine. Contains Suntheanine®, pure L-theanine from Taiyo International, Inc.

Serving Size: 2 Capsules

Servings Per Container: 30

Amount Per Serving:

GABA (Gamma-Aminobutyric acid)

500 mg

L-Theanine

200 mg

Other ingredients: Hydroxypropyl methylcellulose, cellulose, L-leucine, silicon dioxide.

Suggested Use: As a dietary supplement, 1 or 2 capsules daily between meals, or as directed by a healthcare practitioner.

Warning: Contraindicated with drugs or use only under the supervision of a healthcare practitioner.

References

Anderson IM, Mortimore C. Adv Exp Med Biol 1999;467:43-55.
Benton D, Donohoe RT. Public Health Nutr 1999;2:403-9.
Eghbali M, Curmi JP, Birnir B, et al. Nature 1997;388:71-5.
Goddard AW, Mason GF, Almai A, et al. Arch Gen Psychiatry 2001;58:556-61.
He P, Wada S, Watanabe N, et al. J of Food Sci, 2000;65:30-33.
Juneja L, Chu D, Okuba T. Trends Food Sci Tech 1999;10:199-204.
Kakuda T, Nozawa A, Unno T, et al. Biosci Biotechnol Biochem 2000;64:287-93.
Kakuda T, Yanase H, Utsunomiya K, et al. Neurosci Lett 2000;289:189-92.
Kimura R, et al. Chem Pharm Bull (Tokyo) 1986;34:3053-7.
Lombard CB. Med J Aust 2000;173 Suppl:S104-5.
Markus CR, Panhuysen G, Tuiten A, et al. Appetite 1998;31:49-65.
Moore P, Gillin C, et al. Arch Gen Psychiatry 1998;55:534-9.
Paul SM, Marangos PJ, Skolnick P, et al. Encephale 1982;8:131-44.

Paz A, Berry EM. Ann Nutr Metab 1997;41:291-8.
Petty F, Trivedi MH, Fulton M, et al. Biol Psychiatry 1995;38:578-91.
Prasad C. Braz J Med Biol Res 1998;31:1517-27.
Sadzuka Y, Sugiyama T, Sonobe T. Toxicol Lett 2000;114:155-62.
Shiah IS, Yatham LN. Life Sci 1998;63:1289-303.
Sugiyama T, Sadzuka Y. Cancer Lett 1998;133:19-26.
Sugiyama T, Sadzuka Y, Tanaka K, et al. Toxicol Lett 2001;121:89-96.
Sundstrom I, Ashbrook D, Backstrom T. Psychoneuroendocrinology 1997;22:25-38.
Unno T, Suzuki Y, Kakuda T, et al. J Agric Food Chem 1999;47:1593-6.
Verger P, Lagarde D, Batejat D, et al. Physiol Behav 1998;64:317-22.
Vescovi PP, Volpi R, Coiro V. Alcohol 1998;16:325-8.
Wells AS, Read NW, Laugharne JD, et al. Br J Nutr 1998;79:23-30.
Yokogoshi H, Kato Y, et al. Biosci Biotechnol Biochem 1995;59:615-8.
Yokogoshi H, Kobayashi M. Life Sci 1998;62:1065-8.
Yokogoshi H, Mochizuki M, Saitoh K. Biosci Biotechnol Biochem 1998;62:816.



Scientific Bio-Logics®

16612 Burke Lane * Huntington Beach
California * 92647 * E-mail: info@sblogics.com
Phone: 714-847=9355 or Fax: 714-848-8311
www.sblogics.com

*THESE STATEMENTS HAVE NOT BEEN EVALUATED BY THE FOOD AND DRUG ADMINISTRATION. THIS PRODUCT IS NOT INTENDED TO DIAGNOSE, TREAT, CURE, OR PREVENT ANY DISEASE.